


Ch. 6 AP Problems 1 of 2



B) $2.5g C_5H_{12} \cdot \frac{1 mol}{72.096} \cdot \frac{5 CO_2}{1 mol C_5H_{12}} = 0.1734 mol CO_2$

$\left(\frac{785}{760}\right) \cdot V = 0.1734 \cdot 0.0821 \cdot 298$ $V = 4.11 L$



C) $5.00g C_5H_{12} \cdot \frac{1 mol C_5H_{12}}{72.096g} = 6.935 \cdot 10^{-2} mol C_5H_{12}$

$\frac{243 kJ}{6.935 \cdot 10^{-2} mol} = \boxed{3.50 \cdot 10^3 \frac{kJ}{mol}}$

D) $\frac{R_{unk}}{R_p} = \frac{\sqrt{M_p}}{\sqrt{M_{unk}}}$ $R_{unk} = 2R_p$ $2 = \frac{\sqrt{72.096}}{\sqrt{M_{unk}}}$

$M_{unk} = \frac{72.096}{4} = \boxed{18.0g/mol}$ $4 = \frac{72.096}{M_{unk}}$

E)




$$\textcircled{2} \text{ A) } \Delta H = [2\Delta H_{\text{fHCl}}^{\circ}] - [\Delta H_{\text{fOCl}_2}^{\circ} + \Delta H_{\text{fH}_2\text{O}}^{\circ}]$$

$$-46 \text{ kJ} = [2 \cdot -92.3] - [\Delta H_{\text{fOCl}_2}^{\circ} + -242]$$

$$-46 \text{ kJ} = [-184.6] - \Delta H_{\text{fOCl}_2}^{\circ} + 242$$

$$-46 \text{ kJ} = 57.4 - \Delta H_{\text{fOCl}_2}^{\circ}$$

$$\Delta H_{\text{fOCl}_2}^{\circ} = 1.03 \cdot 10^2 \frac{\text{kJ}}{\text{mol}}$$



$$\text{C) } \Delta H_{\text{fOCl}_2}^{\circ} = \left[\frac{1}{2}(\text{O}=\text{O}) + (\text{Cl}-\text{Cl}) \right] - [2(\text{O}-\text{Cl})]$$

$$1.03 \cdot 10^2 = \left[\frac{1}{2}(498) + (243) \right] - 2(\text{O}-\text{Cl})$$

$$1.03 \cdot 10^2 = 492 - 2(\text{O}-\text{Cl})$$

$$-389 = -2(\text{O}-\text{Cl})$$

$$(\text{O}-\text{Cl}) = 195 \frac{\text{kJ}}{\text{mol}}$$

D) Oxygen has a double bond while Cl_2 has only a single bond. Double bonds are stronger and therefore harder to break because there are more e^- being shared between the atoms.

AP[®] CHEMISTRY
2005 SCORING GUIDELINES (Form B)

Question 7

Answer the following questions about thermodynamics.

Substance	Combustion Reaction	Enthalpy of Combustion, ΔH_{comb}° , at 298 K (kJ mol ⁻¹)
H ₂ (g)	H ₂ (g) + $\frac{1}{2}$ O ₂ (g) → H ₂ O(l)	-290
C(s)	C(s) + O ₂ (g) → CO ₂ (g)	-390
CH ₃ OH(l)		-730

- (a) In the empty box in the table above, write a balanced chemical equation for the complete combustion of one mole of CH₃OH(l). Assume products are in their standard states at 298 K. Coefficients do not need to be in whole numbers.

$\text{CH}_3\text{OH}(l) + \frac{3}{2}\text{O}_2(g) \rightarrow \text{CO}_2(g) + 2\text{H}_2\text{O}(l)$	One point is earned for the correct products. One point is earned for balancing the equation.
--	--

- (b) On the basis of your answer to part (a) and the information in the table, determine the enthalpy change for the reaction C(s) + H₂(g) + H₂O(l) → CH₃OH(l).

Adding the following three equations, $\text{C}(s) + \text{O}_2(g) \rightarrow \text{CO}_2(g) \quad -390 \text{ kJ mol}^{-1}$ $\text{H}_2(g) + \frac{1}{2}\text{O}_2(g) \rightarrow \text{H}_2\text{O}(l) \quad -290 \text{ kJ mol}^{-1}$ $\text{CO}_2(g) + 2\text{H}_2\text{O}(l) \rightarrow \text{CH}_3\text{OH}(l) + \frac{3}{2}\text{O}_2(g) \quad +730 \text{ kJ mol}^{-1}$ yields this equation: C(s) + H ₂ (g) + H ₂ O(l) → CH ₃ OH(l) +50 kJ mol ⁻¹	One point is earned for the correct equations. One point is earned for the correct value of ΔH° .
---	---

- (c) Write the balanced chemical equation that shows the reaction that is used to determine the enthalpy of formation for one mole of CH₃OH(l).

$\text{C}(s) + 2\text{H}_2(g) + \frac{1}{2}\text{O}_2(g) \rightarrow \text{CH}_3\text{OH}(l)$	One point is earned for the correct equation.
---	---

AP[®] CHEMISTRY
2005 SCORING GUIDELINES (Form B)

Question 7 (continued)

(d) Predict the sign of ΔS° for the combustion of $\text{H}_2(\text{g})$. Explain your reasoning.

<p>ΔS° for the combustion of $\text{H}_2(\text{g})$ is negative. Both reactants are in the gas phase and the product is in the liquid phase. The liquid phase is much more ordered than the gas phase, so the product is more ordered compared to the reactants, meaning that ΔS° is negative.</p> <p>(Note: There are fewer moles of products than reactants, which also favors a more ordered condition in the products, but the difference in phases is the more important factor.)</p>	<p>One point is earned for the correct sign of ΔS°.</p> <p>One point is earned for a correct explanation.</p>
---	---

(e) On the basis of bond energies, explain why the combustion of $\text{H}_2(\text{g})$ is exothermic.

<p>The combustion of $\text{H}_2(\text{g})$ is exothermic ($\Delta H^\circ < 0$) because more energy is released during the formation of two moles of O–H bonds than is required to break one mole of H–H bonds and one-half of a mole of O–O bonds.</p>	<p>One point is earned for the correct explanation.</p>
---	---